



DIR/Floortime and Autism Spectrum Disorders

Overview and Summary of Scientific and Public Support

Definition of DIR/Floortime

The **D**evelopmental, **I**ndividual Difference, **R**elationship-based (**DIR®/Floortime™**) Model is a developmental, interdisciplinary framework that enables clinicians, parents and educators to construct a comprehensive assessment and intervention program based on the child's and family's unique developmental profile^{1, 2}. It enables the individual working with the child to enter the child's world, bring the child into a shared world, and, from there, interact with the child in ways that build the foundations for social, emotional, and intellectual development. The profile includes understanding the child's emotional, social, and intellectual level; individual differences in motor, sensory, and language functioning; and the existing caregiver, child and family functioning and interaction patterns.^{3 4 5}

Built on years of research in developmental psychology^{6, 7} that underscores the importance of early relationships and family functioning⁸, the DIR/Floortime Model also integrates research contributions from various disciplines, such as speech and language pathology^{9, 10, 11}, occupational therapy^{12, 13, 14}, social work¹⁵, and marriage and family therapy. Neuroscience research lends further support to developmental interventions^{16, 17}. Early relationships and reciprocal interactions between infants and caregivers appear vital for the creation of healthy brain development, including for example wiring the pre-frontal cortex. These diverse contributions provide a rich structure so that clinicians can understand the child and family's unique profile.^{18 19 20} This understanding is crucial for implementing a comprehensive assessment and intervention program for children with autism spectrum disorder (ASD) as well as related special needs conditions²¹.

The objectives of the DIR/Floortime Model are to build the healthy foundations^{22, 23} necessary for a child to develop social, emotional, and intellectual capacities²⁴. As stressed in the general literature that promotes healthy emotional and social development^{25, 26}, two of the key elements to building these foundations are: spontaneous communication²⁷ between children and their caregivers and nurturing relationships²⁸, that promote joyful and pleasant engagement²⁹. These two elements are basic components of the DIR/Floortime Model.

Importantly, the National Research Council of the National Academy of Sciences, in their 2001 landmark report, *Educating Children with Autism*³⁰, called for tailoring the treatment approach to unique features of the individual child and recommended to give priority to interventions that promote functional, spontaneous communication.

Children with ASD differ from one another—in the ways they engage, relate, and communicate and in the ways they respond to sensations and plan and sequence their actions. These differences mean that each child requires an intervention approach tailored to his uniqueness, an intervention that must also consider the home setting. A family where both parents work full-time or where a single parent struggles to put food on the table³¹ has different needs and dynamics than a family where one parent stays home with the child full-time. Thus, each family and team working with the child needs a choice of interventions to formulate a program that optimizes the development for that particular child. The goals of that program, regardless of the approach used, must be to strengthen the child’s core deficits, namely: building the foundations for relating, communicating and thinking. The DIR/Floortime Model is especially beneficial to children with Autism Spectrum Disorders (ASD) and other developmental and/or emotional challenges³²

The DSMIV-TR criteria describe the triad of deficits for children with ASD, which includes qualitative impairments in social interaction and communication as well as restrictive repetitive patterns of behavior, interests, and activities. The DIR/Floortime model provides a roadmap to address these core deficits. It emphasizes the basic foundations of relating, communicating and thinking and builds the core developmental, social, and emotional capacities. Within this framework to build strong foundations, it addresses symptoms such as stereotypical behaviors, self stimulation, and self absorption³³.

Scientific, Professional, Government, and Public Support

The DIR/Floortime Model has been described by the [Centers for Disease Control and Prevention \(CDC\)](#), [Easter Seals](#), and [Autism Speaks](#) as one of the most common interventions for children with ASD. The DIR/Floortime Model has been featured in several news articles, including “[A Tale of Two Schools](#)” (Time, May 2006), and “[Reaching an Autistic Teenager](#)” (New York Times, October 17, 2008). ABC News reported on this latter story with a segment during [Good Morning America](#), emphasizing the importance of social-emotional development to healthy intellectual growth. [ABC News](#) also recently interviewed (April 2009) Erik Linthorst, the producer of the film “Autistic-Like: Graham’s Story”, who describes his [journey from ABA to DIR/Floortime](#) with his son Graham. Elaine Hall, the producer of the Emmy Award winning HBO documentary, [AUTISM: THE MUSICAL](#) has publicly acknowledged how the DIR/Floortime model helped her son and inspired this documentary and [The Miracle Project](#).

Several well conducted studies are showing efficacy of interventions that incorporate a developmental approach, are tailored to the children’s individual abilities, strengthen parent-child interaction and relationships, and focus more on joint attention, social engagement and

reciprocity, and symbolic play, core deficits in ASD^{34, 35, 36, 37, 38, 39, 40, 41}. The DIR/Floortime Model incorporates all the above elements and is widely disseminated by a strong network of well trained DIR/Floortime professionals, all of whom hold clinical licenses or teaching credentials in their own disciplines, and have years of experience in working with infants and young children with ASD and their families. Moreover, thousands of parents have attended DIR/Floortime conferences and workshops since the early 1990s. Training opportunities keep expanding with the DIR Institute Certificate Program and the new distance learning doctoral program, offered by the Interdisciplinary Council on Developmental and Learning Disorders (ICDL)⁴².

The first study to show initial evidence for the DIR/Floortime Model was published in 1997⁴³. Greenspan & Wieder reviewed charts of 200 children who were diagnosed with autistic spectrum disorder, and who were part of a cohort of children seen by the authors over a period of 8 years. All children met the criteria of autism or pervasive developmental disorder not otherwise specified (PDD-NOS) as described in DSM-III-R and DSM-IV, and scored in the autism range on the Childhood Autism Rating Scale⁴⁴. All 200 cases received a comprehensive relationship and developmentally based intervention program for at least two years, under the supervision of Dr. Stanley Greenspan and/or Dr. Serena Wieder. The children ranged in age from 22 months to 4 years, with the majority between 2.5 and 3.5 at the initial evaluation. The goal of the review was to reveal patterns in presenting symptoms, underlying processing difficulties, early development, and response to intervention in order to generate hypotheses for future studies. Based on the findings of this study, Greenspan and Wieder hypothesized that different underlying processing patterns seemed to include a difficulty in connecting affect and sequencing capacities and could be a possible common denominator, suggesting that difficulties with relating and intimacy are often secondary to underlying processing disturbances. They also suggested that the difficulty in engaging in complex purposeful gestural communication could be an early marker, and that contrary to traditional beliefs, a significant number of children may have relatively better functioning in the first year with a regression in the second and third years when these more complex skills are required for social interaction. According to the authors, the chart review suggested that a number of children with autistic spectrum diagnoses are, with an appropriate intervention program, capable of empathy, affective reciprocity, creative thinking, and healthy peer relationships. The authors also concluded that the intervention approach used in these children, focusing on individual differences, developmental level, and affective interaction could be especially promising. Greenspan and Wieder described that after two years of intervention, 58% of treated children showed improvements and no longer met the criteria for ASD.

The results of the 200 case series led them to publish in the year 2000 the full description of the DIR/Floortime Model (ICDL, 2000). In 2005, Greenspan and Wieder published a ten to fifteen-year follow up study (since the start of treatment) of 16 children diagnosed with ASD that were part of the first 200 case series and were part of the 58% of children who showed great improvements⁴⁵. The children were all boys, ranged in age between 12 and 17, with a mean of 13.9 years. All these children had received a comprehensive relationship and developmentally based intervention program, including Floortime at home and DIR consultation, for at least two years (maximum 5), between ages 2-8.5 years old, under the supervision of Dr. Stanley

Greenspan and/or Dr. Serena Wieder. The authors described that after ten to fifteen years since receiving the intervention, these children became empathetic, creative, and reflective adolescents, with healthy peer relationships and solid academic skills. Based on these findings, the authors suggested that some children with ASD can master the core deficits and reach levels of development formerly thought unattainable.

The DIR/Floortime Model also served as the theoretical framework to develop the Greenspan Social-Emotional Growth Chart (SEGC) (Greenspan, 2004)⁴⁶. This norm-referenced surveillance and screening of key social-emotional milestones in infants and children from birth to 42 months of age is now part of the new Bayley Scales Kit of Infant and Early Childhood Development. Published by PsychCorp, the SEGC was field tested on a representative sample of 1,500 infants and young children and it is now offered as a surveillance and screening instrument for Autism Spectrum Disorders, with a sensitivity of 87% and specificity of 90%.

In 2007 Dr. Solomon and colleagues⁴⁷ published an evaluation of The PLAY Project Home Consultation (PPHC), a widely disseminated program that trains parents of children with autism spectrum disorders in the DIR/Floortime model. Sixty-eight children 2 to 6 years old (average 3.7 years) completed an 8–12 month program where parents were encouraged to deliver 15 hours per week of 1:1 interaction. Pre/post ratings of videotapes by blind raters using the Functional Emotional Assessment Scale (FEAS) showed significant increases ($p \leq 0.0001$) in child subscale scores. That is, 45.5 percent of children made good to very good functional developmental progress. Overall parents' satisfaction with PPHC was 90 percent. Despite some limitations, the pilot study of The PLAY Project Home suggests that the model has potential to both enhance developmental progress of young children with autism and to be a cost-effective intervention.

The PLAY model has evolved from a small, university-based, clinical program into a low-cost train-the-trainer model that has the capacity to be disseminated nationally. The state of Michigan Autism Workgroup, convened by the Dept of Health, includes the PLAY Project among the accepted therapies. The PLAY Project, available in a dozen regions of the state, serves hundreds of children in Michigan and is supported by such well-respected agencies as Easter Seals and Mott Children's Health Center Flint. Furthermore, [Easter Seals Crossroads](#) (Easter Seals 2009) has contracted with P.L.A.Y. Project and is beginning to implement this intervention into all of its charter locations. Through the support of a \$1.85 million grant from the National Institute of Mental Health (NIMH), Richard Solomon, MD, is conducting a three-year randomized, controlled, and blinded clinical trial with research-design guidance from Michigan State University, and community-outreach support from Easter Seals. Drawing participants from five Easter Seals autism service locations, the study compares the outcomes of 60 children who participate in The P.L.A.Y. Project with the outcomes of 60 children who receive standard, community interventions, making it the largest study of its kind. Before and after the 12-month intervention, each child is assessed with a battery of tests to measure developmental level, speech and language, sensory-motor profile, and social skills. By training parents to participate in their child's intervention, the program also promises to be cost-effective. The P.L.A.Y. Project costs under \$4,000 per year, in comparison with other interventions that cost \$40,000 to \$60,000 per year.

To further investigate the efficacy of the DIR/Floortime Model, researchers at the [Milton and Ethel Harris Research Initiative](#), at the York University in Canada, are conducting a randomized controlled trial study. The specific aims of this preliminary study are: to assess (1) the efficacy of a 12 months of intensive DIR/Floortime treatment; (2) the magnitude of the gains made by children receiving 24 months of DIR/Floortime treatment; and (3) the neurophysiologic changes that occur as a result of intensive treatment for autism. The study target interventions for children ages 30-51 months old. Of all the changes that the researchers think are occurring in the brain as a result of DIR/Floortime therapy, one of the most important is a shift from ventral to dorsal systems in the Medial Prefrontal Cortex⁴⁸. Ventral systems are more active when performance is emotion-driven and are more susceptible to interference from anxiety and negative emotions, whereas dorsal systems are more active when performance requires cognitive control, such as response inhibition and self-monitoring. The first results were presented at the 2009 Annual ICDL Conference and will be published soon.

A recent article published by Zwaigenbaum and colleagues⁴⁹ highlights the challenges related to early detection, diagnosis, and treatment of ASD in very young children. The authors outline the principles of effective intervention for infants and toddlers with suspected or confirmed diagnosis of ASD, including responsive and sensitive care-taking, enriched language environments using responsive rather than directive interaction styles, environments that provide opportunities for toddlers to take initiative in their learning, and interventions that are individualized and targeted to specific skills. Furthermore, the authors underscore that existing programs for older children “cannot simply be extrapolated” to younger children. Lastly, a comprehensive review article about Autism published in the Lancet journal in 2009⁵⁰ included DIR/Floortime as one of the promising interventions worth considering.

The Interdisciplinary Council on Developmental and Learning Disorders, ICDL

The Interdisciplinary Council on Developmental and Learning Disorders (ICDL) is a non-profit organization founded by Stanley Greenspan, MD, and Serena Wieder, PhD. Its mission is to engage in, develop, conduct, support, and disseminate programs, research, seminars, and publications on the prevention and treatment of emotional and development disorders in infancy and childhood. ICDL has been a pioneer in its work to advance the identification, prevention, and treatment of developmental and learning disorders. Through its research, training and publications, ICDL extends knowledge of developmental processes and provides a framework ([DIR®/Floortime Model](#)) for understanding and improving interventions with infants, children and adults with challenges in relating, communicating and thinking, including autism spectrum disorders.

ICDL’s [Advisory Board](#) consists of nationally and internationally renowned professionals in the field of infant mental health and developmental disorders.

ICDL’s cutting edge publications include, among others, the [CDC/ICDL Collaboration Report on a Framework for Early Identification and Preventive Intervention of Emotional and](#)

[Developmental Challenges](#), the [Clinical Practice Guidelines: Redefining the Standards of Care For Infants, Children, and families with Special Needs](#), [The Journal of Developmental and Learning Disorders](#), and the [Diagnostic Manual for Infancy and Early Childhood \(ICDL-DMIC\)](#).

ICDL was approved by the state of California to start a degree granting institution, the *ICDL Graduate School*, and offer the first [PhD in Infant Mental Health and Developmental Disorders](#).

ICDL's [DIR® Institute has trained a cadre of world class professionals](#) across multiple areas of expertise. It is extending the reach of the DIR®/Floortime™ model as a fundamental component of intervention programs for individuals with difficulties in relating, communicating and thinking, including autism spectrum disorders.

ICDL has successfully offered 3-day [annual conferences](#) since 1997. Thousands of parents and professionals from multiple disciplines, states and countries have attended these conferences, allowing for rich interdisciplinary dialogue with scientists involved in the most innovative research in the field. Over the past eleven years as ICDL has grown, so has its group of creative and innovative parents. It is awe inspiring how many parents want to share their journey, their ideas and their support with one another through different [online parenting resources](#) and [tributes](#) to the DIR/Floortime model.

References

¹ ICDL (2000). *ICDL Clinical Practice Guidelines: redefining the Standards of Care for Infants, Children, and Families with Special Needs*. The Interdisciplinary Council on developmental and Learning Disorders, Bethesda, MD.

² Greenspan S.I. & Wieder S. (1999). A Functional Developmental Approach to Autism Spectrum Disorders. *JASH*, Vol 24, No.3, 147-161.

³ Feuerstein R. (2000). Mediated learning Experience, Instrumental Enrichment and the learning Propensity Assessment Device. In *ICDL Clinical Practice Guidelines: redefining the Standards of Care for Infants, Children, and Families with Special Needs*. The Interdisciplinary Council on developmental and Learning Disorders, Bethesda, MD.

⁴ Wachs H. (2000). Visual-Spatial Thinking. In *ICDL Clinical Practice Guidelines: redefining the Standards of Care for Infants, Children, and Families with Special Needs*. The Interdisciplinary Council on developmental and Learning Disorders, Bethesda, MD.

⁵ Greenspan S.I. (1997). *The Growth of the Mind and the Endangered Origins of Intelligence*. Reading, MA: Addison Wesley Longman.

-
- ⁶ Greenspan S.I. & Shanker (2004). *The First Idea: How symbols, language and intelligence evolved from our primate ancestors to modern humans*. Cambridge, MA: Perseus Books.
- ⁷ Greenspan S.I. & Wieder (1998). *The Child with Special Needs. Encouraging intellectual and emotional growth*. Reading, MA: Perseus Publishing.
- ⁸ Greenspan S.I. (1992). *Infancy and Early Childhood. The practice of clinical assessment and intervention with emotional and developmental challenges*. Madison, CT: International Universities Press.
- ⁹ Wetherby, A.M., & Prizant, B. M. (1995). Facilitating language and communication in autism: Assessment and intervention guidelines. In D. Berkell (ed.), *Autism: Identification, education, and treatment* (pp. 107-133). Hillsdale, NJ: Erlbaum.
- ¹⁰ Tannock, Rosemary, Luigi Girolametto, and Linda Siegal. "Language intervention with children who have developmental delays: Effects of an interactive approach." *American Journal on Mental Retardation* 97, (1992): 145-160.
- ¹¹ Gerber S. & Prizant B. (2000). Speech, Language and Communication Assessment and Intervention for Children. In *ICDL Clinical Practice Guidelines: redefining the Standards of Care for Infants, Children, and Families with Special Needs*. The Interdisciplinary Council on developmental and Learning Disorders, Bethesda, MD.
- ¹² Williamson, G.G., & Anzalone, M.E. (1997). Sensory integration: A key component of the evaluation and treatment of young children with severe difficulties in relating and communicating. *Zero to Three*, 17, 29-36.
- ¹³ Ayres JA. (1979). *Sensory Integration and the Child*. Western Psychological Services. Los Angeles, CA.
- ¹⁴ Case-Smith, J. & Miller, H. (1999). Occupational Therapy with children with pervasive developmental disorders. *American Journal of Occupational Therapy*, 53, 506-513.
- ¹⁵ Shahmoon-Shanok R. (2000). The Action is in the Interaction: Clinical Practice Guidelines for Work with parents of Children with Developmental Disorders. In *ICDL Clinical Practice Guidelines: redefining the Standards of Care for Infants, Children, and Families with Special Needs*. The Interdisciplinary Council on developmental and Learning Disorders, Bethesda, MD.
- ¹⁶ Minshew N.J, & Goldstein, G. (2000). Autism as a Disorder of Complex Information Processing. In *ICDL Clinical Practice Guidelines: redefining the Standards of Care for Infants, Children, and Families with Special Needs*. The Interdisciplinary Council on developmental and Learning Disorders, Bethesda, MD.
- ¹⁷ Mundi, P., Sigman M., Kasari C. (1990). A longitudinal study of joint attention and language development in autistic children. *Journal of Autism and developmental Disorders* 20:115-128.
- ¹⁸ Brazelton, T.B. (1969, 1982). *Infants and Mothers: Differences in Development*. New York: Delacorte Press.
- ¹⁹ Brazelton, T.B. & Nugent, J.K. (1996). *The Neonatal Behavioral Assessment Scale*. 3rd Edition. New York: Cambridge. University Press.

-
- ²⁰ Escalona S. (1968). *The Roots of Individuality*. Chicago: Aldine
- ²¹ Greenspan S.I. & Wieder (2008). *Engaging Autism*. DaCapo Press.
- ²² National Research Council and Institute of Medicine (2000). *From Neurons to Neighborhoods: The Science of Early Childhood Development*. Jack P. Shonkoff and Deborah A. Phillips, eds. Board on Children, Youth, and Families, Commission on behavioral and Social Sciences and Education. Washington, DC: National Academy Press.
- ²³ Greenspan, S.I. (1999). *Building Healthy Minds: The six experiences that create intelligence and emotional growth in babies and young children*. Cambridge, MA: Perseus Books.
- ²⁴ Brazelton T.B., Greenspan, S.I. (2000). *The Irreducible Needs of Children. What Every Child Must Have to Grow, learn, and Flourish*. Perseus Publishing.
- ²⁵ Ainsworth, M., Bell, S.M., & Stayton, D. (1974). Infant-mother attachment and social development: Socialization as a product of reciprocal responsiveness to signals. In M. Richards, ed., *The Integration of the child into a social world*. Cambridge: Cambridge University Press.
- ²⁶ Bowlby, J. (1951). *Maternal care and mental health*. World Health Organization (WHO). Monograph Series, no. 51. Geneva: World Health Organization.
- ²⁷ Gernsbacher M.A., (2006). Toward a behavior of reciprocity. *Journal of Developmental Processes*, 1, 139-152.
- ²⁸ Fogel, A. (1993). *Developing through Relationships*. U of Chicago Press.
- ²⁹ Stern, D. (1974). Mother and infant at play: The dyadic interaction involving facial, vocal, and gaze behaviors. In M. Lewis and L. Rosenblum, eds., *The effect of the infant on its caregiver*. New York: John Wiley & Sons, Inc.
- ³⁰ Lord, Catherine; McGee, James (Editors). Committee on Educational Interventions for Children with Autism. *Educating Children with Autism*. Division of Behavioral and Social Sciences and Education, National Research Council. Washington, DC: National Academy Press (2001).
- ³¹ National Center for Clinical Infant Programs (1987). *Infants in Multirisk Families. Case Studies in Preventive Intervention*. Stanley I. Greenspan, Serena Wieder, Robert A. Nover, Alicia Lieberman, Reginald S. Lourie, Mary E. Robinson, eds. Clinical infant Reports, Number three. International Universities Press.
- ³² Costa G & Witten MR (In press, August 2009). Pervasive Developmental Disorders. In Mowder, Rubinson & Yasik (Eds), *Evidence-Based Practice in Infant and Early Childhood Psychology*. Wiley.
- ³³ Greenspan SI, Brazelton TB, Cordero J, Solomon R, Bauman ML, Robinson R, Shanker S, Breinbauer C. Guidelines to Early Identification, Screening and Clinical Management of Children with Autism Spectrum Disorders. *Pediatrics* 2008 Apr; 121(4):828-30
- ³⁴ Rogers, S. and D. Delalla. (1991). "A comparative study of the effects of a developmentally based instructional model on young children with autism and young children with other disorders of behavior and development." *Topics in Early Childhood Special Education* 11: 29-47.

-
- ³⁵ Mahoney, G., and F. Perales. "Relationship-focused early intervention with children with pervasive developmental disorders and other disabilities: a comparative study." *Journal of Developmental & Behavioral Pediatrics* 26, (2005): 77-85.
- ³⁶ Mahoney, G. & Perales, F. (2003). Using relationship-focused intervention to enhance the social-emotional functioning of young children with autism spectrum disorders. *Topics in Early Childhood Special Education*, 23, 74-86.
- ³⁷ Aldred C, Green J, and Adams C. "A new social communication intervention for children with autism: pilot randomized controlled treatment study suggesting effectiveness." *Journal of Child Psychology & Psychiatry and Allied Disciplines* 45, no. 8 (2004): 1420-30
- ³⁸ Kasari, Connie; Paparella, Tanya; Freeman, Stephanny; Jahromi, Laudan B. "Language outcome in autism: Randomized comparison of joint attention and play interventions." *Journal of Consulting and Clinical Psychology*. Vol 76(1), Feb 2008, 125-137.
- ³⁹ Vismara, L., Colombi, C., Rogers, S. (2009) Can one hour per week of therapy lead to lasting changes in Young Children with autism? *Autism*, 13 (1): 93-115.
- ⁴⁰ Gulsrud, A.C., Jahromi, L.B., Kasari, C. (2010). The Co-Regulation of Emotions Between Mothers and their Children with Autism. *J Autism Dev Disorders*, 40:227-237
- ⁴¹ Kasari C., Gulsrud, A.C., Wong, C., Kwon, S., Locke, J. (2010). Randomized Controlled Caregiver mediated Joint Engagement Intervention for Toddlers with Autism. *J Autism dev Disorders*, published online: 10 February 2010, DOI: 10.1007/s10803-010-0955-5
- ⁴² ICDL (2009a) DIR/Floortime Professionals in United States of America, retrieved from, <http://www.icdl.com/usprograms/clinicians/ListofUSProfessionals.shtml> Accessed July 26, 2009
- ⁴³ Greenspan, S., Wieder, S. (1997) Developmental patterns and outcomes in infants and children with disorders in relating and communicating: A chart review of 200 cases of children with autistic spectrum diagnoses. *Journal of Developmental and Learning Disorders*, 1:87-141.
- ⁴⁴ Western Psychology Services. (1988) Childhood Autism Rating Scale (CARS). Los Angeles, CA.
- ⁴⁵ Greenspan, S.I. and Wieder, S. (2005) Can Children with Autism Master the Core Deficits and Become Empathetic, Creative and Reflective? A Ten to Fifteen Year Follow-up of a Subgroup of Children with Autism Spectrum Disorders (ASD) Who Received a Comprehensive Developmental, Individual-Difference, Relationship-Based (DIR) Approach. *The Journal of Developmental and Learning Disorders*, 9: 39-61.
- ⁴⁶ Greenspan, S.I. (2004). *The Greenspan Social Emotional Growth Chart: A screening questionnaire for infants and young children*. PsychCorp (Harcourt Assessment).
- ⁴⁷ Solomon, R., J. Necheles, C. Ferch, and D. Bruckman. "Pilot study of a parent training program for young children with autism: The P.L.A.Y. Project Home Consultation program." *Autism* 11, no. 3 (2007) 205-224.

⁴⁸ Stieben, J., M. Lewis, I. Granic, P. Zelazo, & D. Pepler, (2007) Neurophysiological mechanism of emotion regulation for subtypes of externalizing children, *Development and Psychopathology*, 19, 455-480.

⁴⁹ Zwaigenbaum L., Bryson S., Lord C., Rogers S., Carter A., Carver L., Chawarska K., Constantino J., Dawson G., Dobkins K., Fein D., Iverson J., Klin A., Landa R., Messinger D., Ozonoff S., Sigman M., Stone W., Tager-Flusberg H., Yirmiya N. (2009) Clinical Assessment and Management of Toddlers With Suspected Autism Spectrum Disorder: Insights From Studies of High-Risk Infants. *Pediatrics*, 123 (5): 1383-1391.

⁵⁰ Levy S.E., Mandell D.S., Schultz, R.T. (2009). Autism. *Lancet*, 374 (9701): 1627-1638.